

FIGHT RETURNING PARKINSON'S SYMPTOMS **NOW**



For the Fighters™

Actor
portrayal

In as early as 10 minutes, you could start to fight Parkinson's symptoms as they return.

Indication

INBRIJA is a prescription medicine used when needed for OFF episodes in adults with Parkinson's treated with regular carbidopa/levodopa medicine. INBRIJA does not replace regular carbidopa/levodopa medicine.

Selected Important Safety Information

Do not use INBRIJA if you take or have taken a nonselective monoamine oxidase inhibitor such as phenelzine or tranylcypromine within the last 2 weeks.

Please see additional Important Safety Information on pages 6-7 and see Patient Information Leaflet and Instructions For Use by visiting www.InbrijaLabel.com.



Inbrija [®]
(levodopa inhalation powder)

Are your Parkinson's symptoms returning?

Many people with Parkinson's disease say they experience symptom return (also known as OFF periods)—and that these returning symptoms are very disruptive.

You may find yourself avoiding going out or making plans because you're concerned about symptom return. This can include missing work or not going out to eat, seeing friends and family, and/or participating in your favorite activities.



Symptom return can take you out of the moment, disrupting your day.



Even when you take your Parkinson's medication regularly, symptoms can still return.

Even one OFF period is one too many.

“The tremors are the worst; they just make functioning very difficult for me.”

—Bettina, real patient with Parkinson's disease

Watch the video



INBRIJA can help address the return of Parkinson's symptoms

INBRIJA is the on-demand levodopa rescue inhaler that can get you back to what you were doing.



INBRIJA starts to work in as early as 10 minutes. In a large study, significant improvement in motor symptoms was seen 30 minutes after taking INBRIJA.



INBRIJA is orally inhaled.

Selected Important Safety Information

Before using INBRIJA, tell your healthcare provider about your medical conditions, including:

- asthma, chronic obstructive pulmonary disease (COPD), or any chronic lung disease
- daytime sleepiness, sleep disorders, sleepiness/drowsiness without warning, or use of medicine that increases sleepiness, including antidepressants or antipsychotics
- dizziness, nausea, sweating, or fainting when standing up
- abnormal movement (dyskinesia)
- mental health problems such as hallucinations or psychosis
- uncontrollable urges like gambling, sexual urges, spending money, or binge eating
- glaucoma
- pregnancy or plans to become pregnant. It is unknown if INBRIJA will harm an unborn baby.
- breastfeeding or plans to breastfeed. Levodopa can pass into breastmilk and it is unknown if it can harm the baby.

Please see additional Important Safety Information on pages 6-7 and see Patient Information Leaflet and Instructions For Use by visiting www.InbrijaLabel.com.

Inbrija[®]
(levodopa inhalation powder)
42 mg capsules

INBRIJA is used as needed



INBRIJA is lightweight and portable
and is easy to carry with you.



Take INBRIJA as prescribed when your Parkinson's symptoms start to return (1 dose per OFF period, no more than 5 doses per day).

PLEASE SEE INSTRUCTIONS FOR USE BY VISITING
www.InbrijaLabel.com.

“I use INBRIJA
anywhere I
want to use it.”

—Jimmy, real patient taking INBRIJA

Watch the video



Selected Important Safety Information

Tell your healthcare provider if you take:

- MAO-B inhibitors
- dopamine (D2) antagonists (including phenothiazines, butyrophenones, risperidone, metoclopramide)
- isoniazid
- iron salts or multivitamins that contain iron salts

Nurse Educators are here to help

If you would like to speak to **Nurse Educators** for additional training on how to use INBRIJA, or if you need assistance with understanding your insurance coverage, Prescription Support Services can assist.



Call toll-free:

1-888-887-3447

8 AM to 8 PM ET,
Monday through Friday

or visit **INBRIJA.com**



Do not drive, operate machinery, or do other activities until you know how INBRIJA affects you. Sleepiness and falling asleep suddenly can happen as late as a year after treatment is started.

Please see additional Important Safety Information on pages 6-7 and see Patient Information Leaflet and Instructions For Use by visiting www.InbrijaLabel.com.

Inbrija[®]
(levodopa inhalation powder)
42 mg capsules

Make sure your doctor knows if you are having OFF periods

Your doctor wants to help. That's why you should be as honest as possible with them.

How are your symptoms impacting your life?

1. Are you ever surprised by symptom return?

- Always Sometimes Never

2. How does the return of your symptoms make you feel?

3. Have your symptoms returned during any of the following situations?

(check all that apply)

- When spending time with family
 While out in public (for example, in a restaurant or park)
 While at work
 On an important day (for example, a wedding or birthday)
 While joining in outdoor activities

4. Have you avoided activities due to concern about symptom return?

- There are many things I won't do any more
 I've stopped doing some things but still enjoy most of my favorite activities
 I'm worried I will have to stop some activities in the future for this reason
 I haven't stopped any activities



Scan the QR code for more helpful information to guide the conversation with your doctor.

Indication

INBRIJA is a prescription medicine used when needed for OFF episodes in adults with Parkinson's treated with regular carbidopa/levodopa medicine. INBRIJA does not replace regular carbidopa/levodopa medicine.

Selected Important Safety Information

Do not use INBRIJA if you take or have taken a nonselective monoamine oxidase inhibitor such as phenelzine or tranylcypromine within the last 2 weeks.

Selected Important Safety Information

Before using INBRIJA, tell your healthcare provider about your medical conditions, including:

- asthma, chronic obstructive pulmonary disease (COPD), or any chronic lung disease
- daytime sleepiness, sleep disorders, sleepiness/drowsiness without warning, or use of medicine that increases sleepiness, including antidepressants or antipsychotics
- dizziness, nausea, sweating, or fainting when standing up
- abnormal movement (dyskinesia)
- mental health problems such as hallucinations or psychosis
- uncontrollable urges like gambling, sexual urges, spending money, or binge eating
- glaucoma
- pregnancy or plans to become pregnant. It is unknown if INBRIJA will harm an unborn baby.
- breastfeeding or plans to breastfeed. Levodopa can pass into breastmilk and it is unknown if it can harm the baby.

Tell your healthcare provider if you take:

- MAO-B inhibitors
- dopamine (D2) antagonists (including phenothiazines, butyrophenones, risperidone, metoclopramide)
- isoniazid
- iron salts or multivitamins that contain iron salts

Do not drive, operate machinery, or do other activities until you know how INBRIJA affects you. Sleepiness and falling asleep suddenly can happen as late as a year after treatment is started.

Tell your healthcare provider if you experience the following side effects:

- **falling asleep during normal daily activities** with or without warning. If you become drowsy, do not drive or do activities where you need to be alert. Chances of falling asleep during normal activities increases if you take medicine that cause sleepiness.
- **withdrawal-emergent hyperpyrexia and confusion** (fever, stiff muscles, or changes in breathing and heartbeat) if you suddenly stop using INBRIJA or carbidopa/levodopa, or suddenly lower your dose of carbidopa/levodopa.
- **low blood pressure** when standing up (that may be with dizziness, fainting, nausea, and sweating). Get up slowly after sitting/lying down.
- **hallucinations and other psychosis** – INBRIJA may cause or worsen seeing/hearing/believing things that are not real; confusion, disorientation, or disorganized thinking; trouble sleeping; dreaming a lot; being overly suspicious or feeling people want to harm you; acting aggressive; and feeling agitated/restless.
- **unusual uncontrollable urges** such as gambling, binge eating, shopping, and sexual urges has occurred in some people using medicine like INBRIJA.
- **uncontrolled, sudden body movements (dyskinesia)** may be caused or worsened by INBRIJA. INBRIJA may need to be stopped or other Parkinson's medicines may need to be changed.
- **bronchospasm** – people with asthma, COPD, or other lung diseases may wheeze or have difficulty breathing after inhaling INBRIJA. If this occurs, stop taking INBRIJA and seek immediate medical attention.
- **increased eye pressure** in patients with glaucoma. Your healthcare provider should monitor this.
- **changes in certain lab values** including liver tests

The most common side effects of INBRIJA are cough, upper respiratory tract infection, nausea, and change in the color of saliva or spit.

Do not orally inhale more than 1 dose (2 capsules) for any OFF period. Do not take more than 5 doses (10 capsules) in a day.

Please see Patient Information Leaflet and Instructions For Use by visiting www.inbrijalabel.com.

Inbrija[®]
(levodopa inhalation powder)
42 mg capsules

Important Safety Information

Do not use INBRIJA if you take or have taken a nonselective monoamine oxidase inhibitor such as phenelzine or tranylcypromine within the last 2 weeks.

Before using INBRIJA, tell your healthcare provider about your medical conditions, including:

- asthma, chronic obstructive pulmonary disease (COPD), or any chronic lung disease
- daytime sleepiness, sleep disorders, sleepiness/drowsiness without warning, or use of medicine that increases sleepiness, including antidepressants or antipsychotics
- dizziness, nausea, sweating, or fainting when standing up
- abnormal movement (dyskinesia)
- mental health problems such as hallucinations or psychosis
- uncontrollable urges like gambling, sexual urges, spending money, or binge eating
- glaucoma
- pregnancy or plans to become pregnant. It is unknown if INBRIJA will harm an unborn baby.
- breastfeeding or plans to breastfeed. Levodopa can pass into breastmilk and it is unknown if it can harm the baby.

Tell your healthcare provider if you take:

- MAO-B inhibitors
- dopamine (D2) antagonists (including phenothiazines, butyrophenones, risperidone, metoclopramide)
- isoniazid
- iron salts or multivitamins that contain iron salts

Do not drive, operate machinery, or do other activities until you know how INBRIJA affects you. Sleepiness and falling asleep suddenly can happen as late as a year after treatment is started.

Please see Patient Information Leaflet and Instructions For Use by visiting www.InbrijaLabel.com.

Tell your healthcare provider if you experience the following side effects:

- **falling asleep during normal daily activities** with or without warning. If you become drowsy, do not drive or do activities where you need to be alert. Chances of falling asleep during normal activities increases if you take medicine that cause sleepiness.
- **withdrawal-emergent hyperpyrexia and confusion** (fever, stiff muscles, or changes in breathing and heartbeat) if you suddenly stop using INBRIJA or carbidopa/levodopa, or suddenly lower your dose of carbidopa/levodopa.
- **low blood pressure** when standing up (that may be with dizziness, fainting, nausea, and sweating). Get up slowly after sitting/lying down.
- **hallucinations and other psychosis** – INBRIJA may cause or worsen seeing/hearing/believing things that are not real; confusion, disorientation, or disorganized thinking; trouble sleeping; dreaming a lot; being overly suspicious or feeling people want to harm you; acting aggressive; and feeling agitated/restless.
- **unusual uncontrollable urges** such as gambling, binge eating, shopping, and sexual urges has occurred in some people using medicine like INBRIJA.
- **uncontrolled, sudden body movements (dyskinesia)** may be caused or worsened by INBRIJA. INBRIJA may need to be stopped or other Parkinson's medicines may need to be changed.
- **bronchospasm** – people with asthma, COPD, or other lung diseases may wheeze or have difficulty breathing after inhaling INBRIJA. If this occurs, stop taking INBRIJA and seek immediate medical attention.
- **increased eye pressure** in patients with glaucoma. Your healthcare provider should monitor this.
- **changes in certain lab values** including liver tests

The most common side effects of INBRIJA are cough, upper respiratory tract infection, nausea, and change in the color of saliva or spit.

Do not orally inhale more than 1 dose (2 capsules) for any OFF period. Do not take more than 5 doses (10 capsules) in a day.

Inbrija [®]
(levodopa inhalation powder)
42 mg capsules

Actor
portrayal

Talk to your doctor about **INBRIJA**



Tell your doctor if the return of Parkinson's symptoms interrupts **YOUR** day and ask if **INBRIJA** is right for you.

In as early as **10 minutes**, you could start to fight Parkinson's disease symptoms as they return.

“ You can advocate for yourself, find the medication that works for you. With **INBRIJA**, I can go out now for a longer time. I can take pictures. I can do stuff I couldn't do. ”

—Bettina, real patient with Parkinson's disease

Watch the video



Hear more patient and carepartner stories at
INBRIJA.com

Selected Important Safety Information

The most common side effects of **INBRIJA** are cough, upper respiratory tract infection, nausea, and change in the color of saliva or spit.

Please see additional Important Safety Information on pages 6-7 and see Patient Information Leaflet and Instructions For Use by visiting www.InbrijaLabel.com.

ACORDA[®]
T H E R A P E U T I C S