

FIGHT RETURNING PARKINSON'S SYMPTOMS NOW



For the Fighters™

Actor
portrayal

In as early as 10 minutes, you could start to fight Parkinson's symptoms as they return.

INBRIJA® (levodopa inhalation powder)

IMPORTANT CONSUMER SAFETY INFORMATION

What is INBRIJA?

INBRIJA is an inhaled levodopa prescription medicine used to treat the return of Parkinson's symptoms (known as OFF episodes) in people with Parkinson's disease who are treated with carbidopa/levodopa medicines. INBRIJA does not replace regular carbidopa/levodopa medicine. It is not known if INBRIJA is safe or effective in children.

Do not use INBRIJA if you: take or have taken a nonselective monoamine oxidase inhibitor such as phenelzine or tranylcypromine within the last 2 weeks.

Please see additional Important Safety Information throughout and see Patient Information Leaflet and Instructions For Use at www.INBRIJAFULLPI.com.



Inbrija [®]
(levodopa inhalation powder)
42 mg capsules

Are your Parkinson's symptoms returning?

Many people with Parkinson's disease say they experience symptom return (also known as OFF episodes)—and that these returning symptoms are very disruptive.

You may find yourself avoiding going out or making plans because you're concerned about symptom return. This can include missing work or not going out to eat, seeing friends and family, and/or participating in your favorite activities.



Symptom return can take you out of the moment, disrupting your day.



Even when you take your oral Parkinson's medication regularly, symptoms can still return.

“The tremors are the worst; they just make functioning very difficult for me.”

—Bettina, real patient with Parkinson's disease

INBRIJA can help address the return of Parkinson's symptoms

INBRIJA is the on-demand levodopa inhaler that can get you back to what you were doing.



INBRIJA may start to work in as early as 10 minutes. In a large study, significant improvement in motor symptoms was seen 30 minutes after taking INBRIJA.



INBRIJA is orally inhaled.



IMPORTANT CONSUMER SAFETY INFORMATION (Continued)

Before using INBRIJA, tell your healthcare provider about your medical conditions, including if you:

- have asthma, chronic obstructive pulmonary disease (COPD), or any chronic lung disease
- have daytime sleepiness, sleep disorders, sleepiness/drowsiness without warning, or use medicine that increases sleepiness, including antidepressants or antipsychotics
- have dizziness, nausea, sweating, or fainting when standing up
- have abnormal movement (dyskinesia)
- have mental health problems such as hallucinations or psychosis
- have uncontrollable urges like gambling, sexual urges, spending money, or binge eating
- have glaucoma
- are pregnant or plan to become pregnant. It is unknown if INBRIJA can harm your unborn baby.
- are breastfeeding or plan to breastfeed. Levodopa can pass into breastmilk and it is unknown if it can harm the baby.

Please see additional Important Safety Information throughout and see Patient Information Leaflet and Instructions For Use at www.INBRIJAFULLPI.com.

Inbrija[®]
(levodopa inhalation powder)

INBRIJA is used as needed



INBRIJA is lightweight and portable
and is easy to carry with you.



Take INBRIJA as prescribed when your Parkinson's symptoms start to return
(1 dose per OFF episode, no more than 5 doses per day). (1 dose is 2 capsules)

PLEASE SEE INSTRUCTIONS FOR USE AT www.INBRIJA.com.

“Whenever I need it morning, noon, or
night INBRIJA is there to help me when
my Parkinson's symptoms return.”

—Bettina, real patient taking INBRIJA

IMPORTANT CONSUMER SAFETY INFORMATION (Continued)

Tell your healthcare provider about all of the medicines you take, including prescription and over-the-counter medicines, vitamins and herbal supplements.

Using INBRIJA and certain other medicines may affect each other and cause serious side effects. **Especially tell your healthcare provider if you take:**

- MAO-B inhibitors
- dopamine (D2) antagonists (including phenothiazines, butyrophenones, risperidone, metoclopramide)
- isoniazid
- iron salts or multivitamins that contain iron salts

Nurse Educators are here to help*

If you would like to speak to **Nurse Educators** for additional training on how to use INBRIJA, or if you need assistance with understanding your insurance coverage, Patient Support Services can assist.



*Paid by Merz Therapeutics. Not affiliated with INBRIJA prescribers.



Call toll-free:

1-833-INBRIJA
(1-833-462-7452)

8 AM to 8 PM ET,
Monday through Friday

or visit **INBRIJA.com**

How should I use INBRIJA?

INBRIJA is **for oral inhalation use only**. Do not swallow or open INBRIJA capsules. Only use INBRIJA capsules with the INBRIJA inhaler. Do not use the INBRIJA inhaler to take any other medicine.

Do not orally inhale more than 1 dose (2 capsules) for any OFF period. Do not take more than 5 doses (10 capsules) in a day.

Please see additional Important Safety Information
on page 6 and see Patient Information Leaflet and
Instructions For Use at www.INBRIJAFULLPI.com.

Inbrija®
(levodopa inhalation powder)

Important Consumer Safety Information (Continued)

What should I avoid while taking INBRIJA?

Do not drive, operate machinery, or do other activities until you know how INBRIJA affects you. Sleepiness and falling asleep suddenly can happen as late as a year after treatment is started.

What are the possible Side Effects of INBRIJA?

INBRIJA can cause serious side effects including:

- **falling asleep during normal daily activities** with or without warning. If you become drowsy, do not drive or do activities where you need to be alert for your safety or the safety of others. Chances of falling asleep during normal activities increases if you take medicine that cause drowsiness.
- **withdrawal-emergent hyperpyrexia and confusion** (fever, stiff muscles, or changes in breathing and heartbeat) if you suddenly stop using INBRIJA or carbidopa/levodopa, or suddenly lower your dose of carbidopa/levodopa.
- **low blood pressure when standing up** (that may or may not happen with dizziness, fainting, nausea, and sweating). Get up slowly after sitting/lying down.
- **hallucinations and other psychosis** - INBRIJA may cause or worsen seeing/hearing/believing things that are not real; confusion, disorientation, or disorganized thinking; trouble sleeping; dreaming a lot; being overly suspicious or feeling people want to harm you; acting aggressive; and feeling agitated/restless.
- **unusual uncontrollable urges** such as gambling, binge eating, shopping, and sexual urges has occurred in some people using medicine like INBRIJA.
- **uncontrolled, sudden body movements** (dyskinesia) may be caused or worsened by INBRIJA. INBRIJA may need to be stopped or other Parkinson's medicines may need to be changed.
- **bronchospasm** - people with asthma, COPD, or other lung diseases may wheeze or have difficulty breathing after inhaling INBRIJA. If this occurs, stop taking INBRIJA and seek immediate medical attention.
- **increased eye pressure** in patients with glaucoma. Your healthcare provider should monitor this.
- **changes in certain lab values** including liver tests.

The most common side effects of INBRIJA are cough, upper respiratory tract infection, nausea, and change in the color of saliva or spit. Another side effect of INBRIJA is sensation of choking right after use.

These are not all the possible side effects of INBRIJA

- Call your doctor for medical advice about side effects.
- You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

The risk information provided here is not comprehensive. To learn more:

- Talk to your health care provider or pharmacist.
- Visit www.inbrija.com to obtain the [Full Prescribing Information, Patient Information, and Instructions for Use](#).
- Call 1-800-367-5109

Make sure your doctor knows if you are having OFF episodes

Your doctor wants to help. That's why you should be as honest as possible with them.

How are your symptoms impacting your life?

1. Are you ever surprised by symptom return?

- ☐ Always ☐ Sometimes ☐ Never

2. How does the return of your symptoms make you feel?

3. Have your symptoms returned during any of the following situations? (check all that apply)

- ☐ When spending time with family
- ☐ While out in public (for example, in a restaurant or park)
- ☐ While at work
- ☐ On an important day (for example, a wedding or birthday)
- ☐ While joining in outdoor activities

4. Have you avoided activities due to concern about symptom return?

- ☐ There are many things I won't do any more
- ☐ I've stopped doing some things but still enjoy most of my favorite activities
- ☐ I'm worried I will have to stop some activities in the future for this reason
- ☐ I haven't stopped any activities

What is INBRIJA?

INBRIJA is an inhaled levodopa prescription medicine used to treat the return of Parkinson's symptoms (known as OFF episodes) in people with Parkinson's disease who are treated with carbidopa/levodopa medicines. INBRIJA does not replace regular carbidopa/levodopa medicine. It is not known if INBRIJA is safe or effective in children.

INBRIJA® (levodopa inhalation powder) IMPORTANT CONSUMER SAFETY INFORMATION

Do not use INBRIJA if you: take or have taken a nonselective monoamine oxidase inhibitor such as phenelzine or tranylcypromine within the last 2 weeks.



Scan the QR code for more helpful information to guide the conversation with your doctor.

Please see additional Important Safety Information throughout and see Patient Information Leaflet and Instructions For Use at www.INBRIJAFULLPI.com.

Inbrija®
(levodopa inhalation powder)

Talk to your doctor about INBRIJA



Tell your doctor if the return of Parkinson's symptoms interrupts YOUR day and ask if INBRIJA is right for you.

In as early as **10 minutes**, you could start to fight Parkinson's disease symptoms as they return.

“ You can advocate for yourself, find the medication that works for you. ”

—Bettina, real patient with Parkinson's disease

Hear more or learn more at **INBRIJA.com**

Important Consumer Safety Information

The most common side effects of INBRIJA are cough, upper respiratory tract infection, nausea, and change in the color of saliva or spit.

Please see additional Important Safety Information throughout and see Patient Information Leaflet and Instructions For Use at www.INBRIJAFULLPI.com.



THERAPEUTICS