

Do you take carbidopa/levodopa and
find that your Parkinson's symptoms still return?

Inbrija [®]
(levodopa inhalation powder)
42 mg capsules

Speak up—your doctor may be able to help

You deserve to be heard about the return of symptoms



**IT'S IMPORTANT FOR YOUR DOCTOR TO FULLY
UNDERSTAND YOUR SYMPTOMS SO THEY CAN
GIVE YOU THE BEST CARE.**

That's why you should be as honest as possible with them. This guide
will help you talk to your doctor about how the return of Parkinson's
symptoms may be impacting you.

INBRIJA is a prescription medicine used when needed for OFF episodes in adults
with Parkinson's treated with regular carbidopa/levodopa medicine. INBRIJA does
not replace regular carbidopa/levodopa medicine.

Selected Important Safety Information

Do not use INBRIJA if you take or have taken a nonselective monoamine oxidase
inhibitor such as phenelzine or tranylcypromine within the last 2 weeks.

**Please see additional Important Safety Information
throughout and the Patient Information Leaflet at
www.INBRIJAlabel.com.**

Think about how your symptoms might impact your life

1 Are you ever surprised by symptom return?

(CHECK ALL THAT APPLY)

- ☐ Always ☐ Never
- ☐ Sometimes

2 How does the return of your symptoms make you feel?

3 Have your symptoms returned during any of the following situations?

(CHECK ALL THAT APPLY)

- ☐ While spending time with family
- ☐ While out in public (for example, in a restaurant or park)
- ☐ While at work
- ☐ On an important day (for example, a wedding or birthday)
- ☐ While joining in outdoor activities

4 Have you avoided any activities due to concern about symptom return?

(CHECK ALL THAT APPLY)

- ☐ There are many things I won't do anymore
- ☐ I'm worried I will have to stop some activities in the future for this reason
- ☐ I've stopped doing some things but still enjoy most of my favorite activities
- ☐ I haven't stopped any activities

Selected Important Safety Information

Before using INBRIJA, tell your healthcare provider about your medical conditions, including:

- asthma, chronic obstructive pulmonary disease (COPD), or any chronic lung disease
- daytime sleepiness, sleep disorders, sleepiness/drowsiness without warning, or use of medicine that increases sleepiness, including antidepressants or antipsychotics
- dizziness, nausea, sweating, or fainting when standing up
- abnormal movement (dyskinesia)
- mental health problems such as hallucinations or psychosis
- uncontrollable urges like gambling, sexual urges, spending money, or binge eating
- glaucoma
- pregnancy or plans to become pregnant. It is unknown if INBRIJA will harm an unborn baby.
- breastfeeding or plans to breastfeed. Levodopa can pass into breastmilk and it is unknown if it can harm the baby.

Inbrija®
(levodopa inhalation powder)
42 mg capsules

Let's talk about when your Parkinson's symptoms return

This is an example of how you can use this guide to start talking with your doctor about symptom return.

"I would like to talk about the ways that symptom return has affected me. I am _____
(YOUR ANSWER TO QUESTION 1)

surprised by symptom return, and it makes me feel _____. These symptoms have
(YOUR ANSWER TO QUESTION 2)

even returned during events in my life like _____.
(YOUR ANSWER TO QUESTION 3)

Because of this, _____. I would like to be able to
(YOUR ANSWER TO QUESTION 4)

take part in these events again. I would like to talk about how we can make that happen."



 **Talk to your doctor to see if
INBRIJA may be right for you.**

“

INBRIJA has allowed me to have Parkinson's and still be able to go out and do all the things that I really want to do.

”

-Jimmy, real INBRIJA patient and clinical trial participant

It's important to make sure your doctor understands how symptom return impacts your life. Be sure to bring this guide with you to your next checkup.

Selected Important Safety Information (cont'd)

Tell your healthcare provider if you take:

- MAO-B inhibitors
- dopamine (D2) antagonists (including phenothiazines, butyrophenones, risperidone, metoclopramide)
- isoniazid
- iron salts or multivitamins that contain iron salts

Please see additional Important Safety Information throughout and the Patient Information Leaflet at www.INBRIJAlabel.com.

Inbrija 
(levodopa inhalation powder)
42 mg capsules

Additional Important Safety Information



Do not drive, operate machinery, or do other activities until you know how INBRIJA affects you. Sleepiness and falling asleep suddenly can happen as late as a year after treatment is started.

Tell your healthcare provider if you experience the following side effects:

- **falling asleep during normal daily activities** with or without warning. If you become drowsy, do not drive or do activities where you need to be alert. Chances of falling asleep during normal activities increases if you take medicine that cause sleepiness.
- **withdrawal-emergent hyperpyrexia and confusion** (fever, stiff muscles, or changes in breathing and heartbeat) if you suddenly stop using INBRIJA or carbidopa/levodopa, or suddenly lower your dose of carbidopa/levodopa.
- **low blood pressure** when standing up (that may be with dizziness, fainting, nausea, and sweating). Get up slowly after sitting/lying down.
- **hallucinations and other psychosis** – INBRIJA may cause or worsen seeing/hearing/believing things that are not real; confusion, disorientation, or disorganized thinking; trouble sleeping; dreaming a lot; being overly suspicious or feeling people want to harm you; acting aggressive; and feeling agitated/restless.
- **unusual uncontrollable urges** such as gambling, binge eating, shopping, and sexual urges has occurred in some people using medicine like INBRIJA.
- **uncontrolled, sudden body movements (dyskinesia)** may be caused or worsened by INBRIJA. INBRIJA may need to be stopped or other Parkinson's medicines may need to be changed.
- **bronchospasm** – people with asthma, COPD, or other lung diseases may wheeze or have difficulty breathing after inhaling INBRIJA. If this occurs, stop taking INBRIJA and seek immediate medical attention.

- **increased eye pressure** in patients with glaucoma. Your healthcare provider should monitor this.
- **changes in certain lab values** including liver tests

The most common side effects of INBRIJA are cough, upper respiratory tract infection, nausea, and change in the color of saliva or spit.

Do not orally inhale more than 1 dose (2 capsules) for any OFF period. Do not take more than 5 doses (10 capsules) in a day.

Please see the Patient Information Leaflet at www.INBRIJAlabel.com.



ACORDA THERAPEUTICS, the stylized ACORDA THERAPEUTICS logo, INBRIJA, and the INBRIJA logo are trademarks of Acorda Therapeutics, Inc.
© 2022 Acorda Therapeutics, Inc. All rights reserved. 11/22 INB13083